




**UNIVERSITI PUTRA MALAYSIA**


**EFFECT OF AVAILABLE SOIL MOISTURE ON THE GROWTH AND  
YIELD OF GROUDNUT (ARACHIS HYPOGAEA L.)**


**MD. ABDUR RAHMAN SARKAR**


**FP 1987 7**

It is hereby certified that we have read this thesis entitled "Effects of Available Soil Moisture on The Growth and Yield of Groundnut (arachis hypogaea L.)" by Md. Abdur Rahman Sarkar, and in our opinion it is satisfactory in terms of scope, quality, and presentation as partial fulfilment of the requirements for the degree of Doctor of Philosophy

  
.....  
ALANG P. ZAINUDDIN, Ph.D.  
Assoc. Professor/Dean of Graduate Studies  
Universiti Pertanian Malaysia  
(Chairman Board of Examiners)


  
.....  
JONATHAN H. WILLIAMS, Ph.D.  
Visiting Scientist  
Department of Agronomy & Soils  
Washington State University  
U.S.A.  
(External Examiner)

  
.....  
MOHAMMAD MD. ALI, Ph.D.  
Associate Professor  
Faculty of Agriculture  
Universiti Pertanian Malaysia  
(Internal Examiner)

  
.....  
LIM ENG SIONG, Ph.D.  
Associate Professor  
Faculty of Agriculture  
Universiti Pertanian Malaysia  
(Internal Examiner/Supervisor)

This thesis was submitted to the Senate of Universiti  
Pertanian Malaysia and was accepted as partial fulfilment of the  
requirements for the degree of Doctor of Philosophy.

Date: 15 DEC 1988

  
.....  
SULAIMAN M. YASSIN, Ph.D.  
Professor,  
Dean of Graduate Studies.

EFFECT OF AVAILABLE SOIL MOISTURE ON THE GROWTH AND  
YIELD OF GROUNDNUT ( Arachis hypogaea L.)

By

MD. ABDUR RAHMAN SARKAR

A thesis submitted in partial fulfilment of the  
requirements for the degree of Doctor of Philosophy  
in the Faculty of Agriculture, University of Agriculture  
Malaysia

June 1987



## DEDICATION

Dedicated to the memory  
of my departed father

## ACKNOWLEDGEMENTS

The author expresses his deep sense of gratitude and indebtedness to his supervisors Dr. Lim Eng Siong and Dr. Mok Chak Kim for their painstaking and scholastic guidance, advice, and constant encouragement during the entire period of this study and for their invaluable constructive criticisms during the preparation of the thesis.

The author owes a debt of gratitude to the Govt. of Malaysia for financing the study through Commonwealth Scholarship.



## TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
TABLE OF CONTENTS	iv
LIST OF TABLES	viii
LIST OF FIGURES	x
LIST OF PLATES	xi
LIST OF APPENDICES	xii
ABSTRACT	xiv
PREFACE	xvi
CHAPTER I. INTRODUCTION	1
CHAPTER II. REVIEW OF LITERATURE	4
Development of Plant Water Deficit	4
Water Stress and the Physiological Processes in Plants	5
Protoplasmic Dehydration	5
Effects on Photosynthesis	6
Effects of Respiration	7
Effects on Transpiration	7
Effects on Translocation	8
Water Stress and the Vegetative Growth of Crop Plants	8
Growth of Leaves	9
Growth of Stems	10



	Page
Growth of Roots and Top-root Ratio	11
Dry Matter Production	12
Recovery of Growth Upon the Alleviation of Stress	13
Effect of Moisture Stress on the Yield of Crops	14
Moisture Stress at the Vegetative Stage and Yield of Crops	15
Moisture Stress at the Reproductive Phase and Yield of Crops	16
Moisture Stress at the Flower Initiation and Flowering Stage and Yield of Crop Plants	16
Moisture Stress and the Stage of Fruit Enlargement	19
Moisture Stress and the Stage of Ripening or Crop Maturity	21
Moisture Stress and Maturity of Crops	22
CHAPTER III. MATERIALS AND METHODS	23
Plant Material	23
Soil	23
Experimental Details	24
Experiment I	26
Experiment II	32
Experiment III	35



	Page
CHAPTER IV. RESULTS	39
Experiment I	39
Vegetative Characters	39
Dry Weight of Haulms	39
Dry Weight of Root	41
Reproductive Characters	41
Production of Flowers	41
Number of Pegs	44
Yield and Yield Component	44
Pod Yield	44
Experiment II	51
Vegetative Growth	51
Length of Stem	51
Dry Weight of Stem	53
Number of Leaves per Plant	53
Leaf Area per Plant	53
Dry Weight of Roots	59
Dry Matter Production (Excluding pods)	59
Reproductive Characters	63
Number of Flowers, Pegs and Pods Produced	63
Pod Development and Yield	65
Quality of Produce	68
Correlation Between Pod Yield and Total Dry Matter Yield	71

	Page
Experiment III	74
Vegetative Growth	74
Dry Weight of Leaves	74
Dry Weight of Stems	74
Number of Branches Produced per Plant	74
Dry Weight of Root	77
Reproductive Characters	77
Production of Flowers	77
Production of Pegs	79
Production of Pods	79
Pod and Seed Size	88
Correlation Between Pod Yield and Total Dry Matter Yield	88
CHAPTER V. DISCUSSION	90
Effect of Available Soil Moisture on Vegetative Growth	90
Effect of Available Soil Moisture on the Reproductive Growth and Yield	93
Important Levels of Available Soil Moisture and Duration of the Moisture Treatment	96
Critical Growth Stage	97
The Limitations and Applicability of Pot Studies to Field Conditions	97
Conclusion	99
CHAPTER VI. SUMMARY AND CONCLUSION	100
BIBLIOGRAPHY	104
APPENDICES	113



## LIST OF TABLES

TABLE	PAGE
I. Effect of the Duration of Moisture Treatment, Growth Stage Treated and Available Soil Moisture on the Dry Weight of Haulm (g/pot)	40
II. Effect of Available Soil Moisture on Root Dry Weight	42
III. Effect of Growth Stage at the Time of Treatment on Root Dry Weight	42
IV. Effect of Moisture Treatment Duration, Growth Stage and Available Soil Moisture on the Mean Number of Flowers Produced per Pot	43
V. Effect of Available Soil Moisture on the Mean Number of Pegs Produced per Pot	46
VI. Effect of Moisture Treatment Duration, Growth Stage and Available Soil Moisture on the Dry Weight of Pods (g/pot)	46
VII. Effect of Moisture Treatment Duration, Growth Stage and Available Soil Moisture on the Number of Mature Pods Produced per Pot	49
VIII. Effect of Moisture Treatment Duration, Growth Stage and Available Soil Moisture on the Dry Weight of Seeds (g/pot)	50
IX. Effect of Available Soil Moisture on Pod Production per Plant	66
X. Effect of Available Soil Moisture on Mature Pod and Seed Yield	69
XI. Effect of Available Soil Moisture on the Number of One, Two and Three-seeded Pods	70
XII. Effect of Growth Stage on Dry Weight of leaves (g/plant)	75
XIII. Effect of Growth Stage at Treatment and Available Soil Moisture on the Dry Weight of Stems (g/plant)	75



ABST	PAGE
IV. Effect of Growth Stage and Available Soil Moisture on the Total Number of Branches Produced per Plant	76
V. Effect of Growth Stage at Treatment on Dry Weight of Roots (g/plant)	78
VI. Effect of Growth Stage at Treatment and Available Soil Moisture on the Total Number of Flowers Produced per Plant	78
VII. Effect of Growth Stage at Treatment and Available Soil Moisture on the Number of Pegs Produced per Plant	81
VIII. Effect of Growth Stage at Treatment and Available Soil Moisture on the Total Number of Pods per Plant	81
IX. Effect of Growth Stage at Treatment and Available Soil Moisture on the Number of Undeveloped Pods per Plant	82
XX. Effect of Available Soil Moisture on the Yield and Yield Components	82
XI. Effect of Growth Stage at Treatment on the Yield and Yield Components	84
XII. Effect of Growth Stage at Treatment and Available Soil Moisture on the Number of Two-seeded Pods per Plant	86
XIII. Effect of Growth Stage at Treatment and Available Soil Moisture on the Number of One-seeded Pods per Plant	87



## LIST OF FIGURES

No	Figure	PAGE
1.	Soil Moisture Characteristic Curve for Serdang Colluvium Soil	25
2.	Planting and Harvesting Dates and Moisture Treatment Period in Experiment I	30
3.	Planting and Harvesting Dates and Moisture Treatment Period in Experiment III	37
4.	Effect of Available Soil Moisture on Main Stem Length	52
5.	Effect of Available Soil Moisture on Dry Weight of Stems	54
6.	Effect of Available Soil Moisture on the Length of First Pair of $n + 1$ Branches	55
7.	Effect of Available Soil Moisture on the Length of Second Pair of $n + 1$ Branches	56
8.	Effect of Available Soil Moisture on Number of Leaves	57
9.	Effect of Available Soil Moisture on Leaf Area	58
10.	Effect of Available Soil Moisture on Dry Weight of Leaves	60
11.	Effect of Available Soil Moisture on Dry Weight of Roots	61
12.	Effect of Available Soil Moisture on Dry Matter Production	62
13.	Effect of Available Soil Moisture on the Number of Flowers, Pegs and Pods	64
14.	Effect of Available Soil Moisture on the Rate of Pod Development	67
15.	Relationship Between Pod Yield and Total Dry Matter (Experiment II.)	72
16.	Relationship Between Pod Yield and Total Dry Matter (Experiment III).	89

## LIST OF PLATES

PLATE	PAGE
I. Method of Weighing the Pots for Watering	29
II. Plants Grown at Different Available Soil Moisture Regimes (Experiment II.)	73



## LIST OF APPENDICES

	PAGE
I. Mean monthly minimum and maximum air temperature (°C) and relative humidity (%) recorded in the glasshouse during the experiments	113
II. Experiment I: Mean square values for the dry weight of haulm and roots	114
III. Experiment I : Mean square values for the number of flowers and pegs produced per pot	115
IV. Experiment I: Mean square values for the yield and its components	116
V. Experiment II: Mean square values for the length of the main stem (cm) measured every 7 days beginning 25 days after planting	117
VI. Experiment II: Mean square values for the dry weight of stems (g/plant) sampled at different days after planting	118
VII. Experiment II: Mean square values for the length of the first pair of n + 1 branches (cm) measured every 7 days beginning 25 days after planting	119
VIII. Experiment II : Mean square values for the length of the second pair of n + 1 branches (cm) measured every 7 days beginning 25 days after planting	120
IX. Experiment II: Mean square values for total number of branches per plant at harvest	121
X. Experiment II: Mean square values for the number of leaves per plant measured at different days after planting	122
XI. Experiment II: Mean square values for the leaf area per plant (cm <sup>2</sup> ) sampled at different days after planting	123
XII. Experiment II: Mean square values for the dry weight of leaves (g/plant) sampled at different days after planting	124



	PAGE
XIII. Experiment II: Mean square values for the dry weight of root (g/plant) sampled at different days after planting	125
XIV. Experiment II: Mean square values for the total dry matter (g/plant) sampled at different days after planting	126
XV. Experiment II: Mean square values for the numbers of flowers, pegs and total pods per plant	127
XVI. Experiment II: Mean square values for the dry weight of developing pods (g/plant) sampled at different days after planting	128
XVII. Experiment II: Mean square values for the yield and its components	129
XVIII. Experiment II: Mean square values for the number of pods per plant (transformed) in various pod categories	130
XIX. Experiment III: Mean square values for the dry weight of leaves, dry weight of stems, total number of branches and dry weight of root per plant	131
XX. Experiment III: Mean square values for the number of flowers, pegs and total pods per plant	132
XXI. Experiment III: Mean square values for the number of pods per plant in various pod categories	133
XXII. Experiment III: Mean square values for the yield and its components	134





An abstract of the thesis presented to the Senate of  
Universiti Pertanian Malaysia in partial fulfilment of the  
requirements for the Degree of Doctor of Philosophy.

EFFECT OF AVAILABLE SOIL MOISTURE ON THE  
GROWTH AND YIELD OF GROUNDNUT (Arachis hypogaea L.)

By

Md. Abdur Rahman Sarkar

Chief Supervisor : Associate Professor Dr. Lim Eng Siong  
Supervisor : Associate Professor Dr. Mok Chak Kim  
Faculty : Agriculture

Experiments were carried out under glasshouse conditions  
with a locally grown Spanish type groundnut variety V 13 to  
determine the effect of available soil moisture on the growth  
and yield.



The vegetative growth was found to be depressed by the prolonged deficit of available soil moisture at early flowering. The pod yield was found to be reduced by moisture deficits of both short and prolonged duration occurring at early flowering and pegging stages in comparison to podding stage. However, the reduction in pod yield was more severe due to the prolonged deficit of available soil moisture at early flowering. Twenty percent available soil moisture reduced pod yield more severely during the prolonged duration treatment.

The vegetative growth (particularly in terms of dry matter) was adversely affected when irrigation was scheduled at or below 70 percent available soil moisture. As a consequence of the adverse effect of low levels of available soil moisture on the vegetative growth, the pod and seed yields were also adversely reduced.

The flowering and podding were found to be more sensitive to low levels of available soil moisture in respect of yield.

With a view to maximizing yield, the groundnut crop should preferably be irrigated to maintain the level of available soil moisture at field capacity especially during the stage of flowering and podding.



Abstrak tesis yang dikemukakan kepada Senat Universiti Pertanian Malaysia sebagai memenuhi sebahagian dari syarat-syarat untuk memperolehi Ijazah Doktor Falsafah.

KESAN KELEMBAPAN TANAH YANG SEDIA ADA KE ATAS TUMBESARAN  
DAN PENGHASILAN KACANG TANAH (Arachis hypogaea L.)

Oleh

Md. Abdur Rahman Sarkar

Ketua Penelia : Profesor Madya Dr. Lim Eng Siong  
Penelia : Profesor Madya Dr. Mok Chak Kim  
Fakulti : Pertanian

Eksperimen telah dijalankan di bawah keadaan rumah kaca ke atas satu varieti kacang tanah Spanish tempatan, varieti V 13, untuk menentukan kesan kelembapan tanah yang sedia ada terhadap tumbesaran dan penghasilannya.

Tumbesaran tampak pada peringkat pembungaan awal telah didapati dirosotkan oleh kekurangan kelembapan tanah yang sedia ada yang berpanjangan. Hasil lenggai juga didapati telah dikurangkan oleh kekurangan kelembapan yang berlaku pada tempoh jangka pendek dan panjang pada peringkat pembungaan awal dan benangan berbanding dengan peringkat pelenggaian. Walau

bagaimanapun, pengurangan hasil lenggai berlaku dengan lebih teruk akibat daripada kekurangan air tanah yang sedia ada yang berpanjangan pada peringkat pembungaan awal. Kelembapan tanah yang sedia ada pada paras 20 peratus sangat menjejaskan hasil lenggai.

Tumbesaran tampang (Khasnya dari segi bahan kering) telah terjejas apabila bekalan air diadakan pada paras kelembapan tanah yang sedia ada pada atau di bawah 70 peratus. Kesan buruk ke atas tumbesaran tampang yang disebabkan oleh kelembapan tanah yang rendah ialah hasil lenggai dan biji benih menjadi kurang dengan teruk.

Dari segi penghasilan, peringkat pembungaan dan pelenggaian telah didapati lebih sensitif kepada paras rendah kelembapan tanah yang sedia ada.

Untuk mendapatkan hasil yang maksimum, tanaman kacang tanah perlu diberikan bekalan air supaya paras kelembapan tanah yang sedia ada dikekalkan pada had basah (field capacity), istimewa sekali pada masa peringkat pembungaan dan pelenggaian.

## CHAPTER I

### INTRODUCTION

Groundnut (Arachis hypogaea L.) which belongs to the family Papilionaceae in the Order Leguminosae, is regarded as one of the leading crops in the world for the production of oil and plant protein. It is extensively cultivated in the tropical and sub-tropical countries for direct use as food, for the oil, and for the protein rich meal produced after oil extraction.

Erratic rainfall, in the tropical and sub-tropical countries is one of the important causes responsible for the low yield in groundnut as it is usually grown there as a rainfed crop. The crop generally requires an evenly distributed rainfall of at least 550 mm (Sellschop, 1966) during the growing season when grown as a rainfed crop. Soil moisture deficits frequently occur under rainfed condition when anticipated rain does not occur in time or when rainfall is scanty. Under such conditions, the crop has to depend upon the stored soil moisture which is soon depleted through evapotranspiration. Consequently the growth and yield of the crop is severely depressed. Prolonged and severe moisture deficits can result in complete crop failure unless the soil moisture is replenished in time. Natural rainfall supplemented with irrigation can ensure favourable soil moisture regimes



conducive to optimum yield. However, irrigation is costly and the quantity of water is usually limited. Therefore, judicious use of irrigation is desirable.

Plants are not equally sensitive to soil moisture deficits at different stages of growth in terms of growth and yield reduction. Rather, there are some specific stages of growth during which soil moisture deficits are more prone to reduce the growth and yield. Scanty rainfall during the moisture sensitive growth stages, can result in severe yield reduction unless depleted available soil moisture is replenished with supplemental irrigation. The irrigation cost may be minimized by supplying adequate water at the moisture-sensitive stage and less water at other stages of growth.

The determination of the level of available soil moisture upto which irrigation can be delayed, is an important cost reducing aspect in groundnut irrigation. The interval between irrigations should be as large as possible without appreciable reduction in crop yield. This will also ensure the application of the right amount of water. Over-irrigation, which is costly and detrimental to the crop, can thus be avoided.

Groundnut is one of the promising short-term cash crops in Malaysia included in the crop diversification programme. It is generally grown here as a rainfed crop. It is, therefore, probable that the groundnut yield can be reduced in times of

inadequate rainfall). There is limited and sporadic information on the growth and yield of groundnut in relation to soil moisture deficits under local condition. Research work in this field thus warrants due attention.

The present study was, therefore, carried out with the following objectives :

1. . To determine the growth and yield response of groundnut at different growth stages to short and prolonged duration at various levels of available soil moisture.
2. To determine the optimum level of available soil moisture for the growth and production of groundnut, and to determine the moisture sensitive growth stage.

## CHAPTER II

### REVIEW OF LITERATURE

This review of literature covers the development of plant water deficit and its effects on physiological processes as well as on the growth, development and yield of crop plants in general and groundnut in particular.

#### Development of Plant Water Deficit

Plant water deficits generally occur when water loss through transpiration is more than water absorption. According to Kramer (1963), water deficits in plants can be caused either by excessive loss of water or by inadequate absorption, or by a combination of the two. The water potential in a non-transpiring plant approaches the water potential in the soil, but with the commencement of transpiration, a gradient in water potential develops from the soil, to and through the plant; thus water moves from the soil, through the plant, and out to the atmosphere (Crischer and Hagan, 1965). Temporary midday deficits occur in rapidly transpiring plants because the resistance to water movement through roots causes absorption to lag behind transpiration even in moist soil. Longer-term and more severe water deficits develop when decrease



cause decreased absorption of water. Thus, daily cycles in water stress are controlled chiefly by transpiration, but long term, severe water deficits develop chiefly because of decreasing availability of soil water (Kramer, 1969). The decreasing availability of soil water causes an increase in the water potential gradient between the soil and the root. As a result, a greater soil-root water potential difference is required to move water from soil to roots to replace the loss by transpiration. At this stage the rate of water movement from the soil to the roots is so slow that recovery from wilting is not possible and permanent wilting occurs.

#### Water Stress and the Physiological Processes in Plants

Water stress affects physiological processes such as protoplasmic dehydration, photosynthesis, respiration, transpiration and translocation of photosynthates.

#### Protoplasmic Dehydration

The dehydration of the protoplasm has profound effect on the various physiological processes in plant life. According to Stocker (1960), the dehydration of the protoplasm occurs in two stages - the reaction phase, when plants are first subjected to water stress and the restitution and hardening phase, which occurs if water stress lasts for several days. The reaction phase is characterized by a decrease in viscosity of the protoplasm, increased permeability to water, urea and glycerine, increased proteolysis, and increased respiration. In the restitution phase,